

Application/Control Number: 09/374,704

Page 2

Art Unit: 1635

CPTO

K.T.

11/22/04

Art Unit: 1635

1. (Twice Amended) A polyamide molecule that specifically binds to base pairs in the minor groove of a DNA molecule, said polyamide molecule comprising:

one or more amino acids comprising a moiety selected from the group consisting of N-methylpyrrole, 3-hydroxy-N-methylpyrrole, and N-methylimidazole, wherein one or more of said amino acid(s) are not α -amino acids; and

a positive patch consisting of a rigid group adjacent to a positively charged group, said rigid group comprising a first and a second amino acid; said first amino acid being selected from the group consisting of arginine, proline, lysine, and hydroxyproline; and said second amino acid being selected from the group consisting of proline, glycine, serine, threonine, leucine, isoleucine, valine, alanine, and hydroxyproline.

3. (Amended) The polyamide of claim 2 wherein said first amino acid is arginine and said second amino acid is proline.

4. (Amended) The polyamide of claim 1 wherein the positively charged group comprises a synthetic or naturally occurring amino acid having a net positive charge.

5. (Amended) The polyamide of claim 1 wherein said positively charged group is selected from the group consisting of a primary amino group, secondary amino group, tertiary amino group, quaternary amino group, guanidinium group, and an amidinium group.

6. (Twice amended) The polyamide of claim 1 wherein said positively charged group is selected from the group consisting of arginine, lysine, and histidine.

Art Unit: 1635

7. (Amended) The polyamide of claim 1 wherein said positively charged group is arginine.


8. (Amended) The polyamide of claim 1 wherein the positive patch comprises the amino acid sequence Arg-Pro-Arg.

9. (Amended) The polyamide of claim 1 wherein the polyamide has three or four carboxamide binding pairs.

10. (Amended) The polyamide of claim 1 wherein the polyamide comprises an (R)-2,4-diaminobutyric acid hairpin turn that facilitates specific binding to base pairs in the minor groove of a DNA molecule.

11. (Amended) The polyamide of claim 10 wherein the R-2-amino group is derivatized to form an acid amide.

12. (Twice Amended) The polyamide of claim 1 having the formula:



wherein γ is $-\text{NH}-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CONH}-$ hairpin linkage derived from γ -aminobutyric acid or a chiral hairpin linkage derived from 2,4-diaminobutyric acid;

X_1/X_6 , X_2/X_5 , and X_3/X_4 represent three carboxamide binding pairs which bind DNA base pairs and are selected from the group consisting of 3-hydroxy-N-methylpyrrole/N-methylpyrrole (Hp/Py), N-methylpyrrole/3-hydroxy-N-methylpyrrole (Py/Hp), N-methylpyrrole/N-methylimidazole (Py/Im), N-methylimidazole/N-methylimidazole (Im/Py), and N-methylpyrrole/N-methylpyrrole (Py/Py) to correspond to the DNA base pair in the minor groove to be bound; and

A represents said positive patch consisting of a rigid group adjacent to a positively charged group.

13. (Amended) The polyamide of claim 12 wherein the positive patch comprises the amino acid sequence Arg-Pro-Arg.

PATENT

14. (Twice amended) The polyamide of claim 1 having the formula:



wherein γ is $-\text{NH}-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CONH}-$ hairpin linkage derived from γ -aminobutyric acid or a chiral hairpin linkage derived from 2,4-diaminobutyric acid;

X_1/X_8 , X_2/X_7 , X_3/X_6 , and X_4/X_5 represent four carboxamide binding pairs which bind DNA base pairs and are selected from the group consisting of Hp/Py, Py/Hp, Py/Im, Im/Py, and Py/Py to correspond to the DNA base pair in the minor groove to be bound; and

A represents said positive patch consisting of a rigid group adjacent to a positively charged group.

15. (Amended) The A polyamide of claim 14 wherein the positive patch comprises the amino acid sequence Arg-Pro-Arg.

16. (Twice amended) The polyamide of claim 1 having the formula:




wherein γ is $-\text{NH}-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CONH}-$ hairpin linkage derived from γ -aminobutyric acid or a chiral hairpin linkage derived from 2,4-diaminobutyric acid;

X_1/X_{10} , X_2/X_9 , X_3/X_8 , X_4/X_7 , and X_5/X_6 represent five carboxamide binding pairs which bind DNA base pairs and are selected from the group consisting of Hp/Py, Py/Hp, Py/Im, Im/Py, and Py/Py to correspond to the DNA base pair in the minor groove to be bound; and

A represents said positive patch consisting of a rigid group adjacent to a positively charged group.

17. (Amended) The polyamide of claim 16 wherein the positive patch comprises the amino acid sequence Arg-Pro-Arg.

18. (Twice amended) The polyamide of claim 1 having the formula:



wherein γ is $-\text{NH}-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CONH}-$ hairpin linkage derived from γ -aminobutyric acid or a chiral hairpin linkage derived from 2,4-diaminobutyric acid;


X_1/X_{12} , X_2/X_{11} , X_3/X_{10} , X_4/X_9 , X_5/X_8 , and X_6/X_7 represent six carboxamide binding pairs which bind DNA base pairs and are selected from the group consisting of Hp/Py, Py/Hp, Py/Im, Im/Py, and Py/Py to correspond to the DNA base pair in the minor groove to be bound; and

A represents said positive patch consisting of a rigid group adjacent to a positively charged group.

19. (Amended) The polyamide of claim 18 wherein the positive patch comprises the amino acid sequence Arg-Pro-Arg.

Art Unit: 1635

Claim 20. (Amended) A tandem-linked polyamide having the formula:



wherein γ is $-\text{NH}-\text{CH}_2-\text{CH}_2-\text{CONH}-$ hairpin linkage derived from γ -aminobutyric acid or a chiral hairpin linkage derived from 2,4-diaminobutyric acid;


X_1/X_6 , X_2/X_5 , X_3/X_4 , X'_1/X'_6 , X'_2/X'_5 , and X'_3/X'_4 represent carboxamide binding pairs which bind DNA base pairs and are selected from the group consisting of Hp/Py, Py/Hp, Py/Im, Im/Py, and Py/Py to correspond to the DNA base pair in the minor groove to be bound;

L represents an amino acid linking group selected from the group consisting of β -alanine and 5-aminovaleric acid (δ);

P represents a polyamide selected from the group consisting of $X_1X_2X_3X_4X_5X_6$, $X_1X_2X_3X_4X_5X_6X_7X_8$, $X_1X_2X_3X_4X_5X_6X_7X_8X_9X_{10}$, and $X_1X_2X_3X_4X_5X_6X_7X_8X_9X_{10}X_{11}X_{12}$, where X_1 - X_{12} are independently selected from the group consisting of β -alanine, pyrrole, hydroxypyrrole and imidazole; and

A represents a positive patch consisting of a rigid group adjacent to a positively charged group [such that a positive charge is delivered to the phosphate backbone or the major groove of a DNA molecule].

Claim 21. (Amended) A tandem-linked polyamide having the formula:



wherein γ is $-\text{NH}-\text{CH}_2-\text{CH}_2-\text{CONH}-$ hairpin linkage derived from γ aminobutyric acid or a chiral hairpin linkage derived from 2,4-diaminobutyric acid;

X_1/X_6 , X_2/X_7 , X_3/X_6 , X_4/X_5 , X'_1/X'_6 , X'_2/X'_7 , X'_3/X'_6 , and X'_4/X'_5 , represent carboxamide binding pairs which bind DNA base pairs and are selected from the group consisting of Hp/Py, Py/Hp, Py/Im, Im/Py, and Py/Py to correspond to the DNA base pair in the minor groove to be bound;


L represents an amino acid linking group selected from the group consisting of β -alanine and 5-aminovaleric acid (δ);

P represents a polyamide selected from the group consisting of $X_1X_2X_3X_4X_5X_6$, $X_1X_2X_3X_4X_5X_6X_7X_8$, $X_1X_2X_3X_4X_5X_6X_7X_8X_9X_{10}$, and $X_1X_2X_3X_4X_5X_6X_7X_8X_9X_{10}X_{11}X_{12}$, where X_1 - X_{12} are independently selected from the group consisting of β -alanine, pyrrole, hydroxypyrrole and imidazole; and

Art Unit: 1635

A represents a positive patch consisting of a rigid group adjacent to a positively charged group [such that a positive charge is delivered to the phosphate backbone or the major groove of a DNA molecule].

22. (Twice Amended) A tandem-linked polyamide having the formula:



wherein γ is $-\text{NH}-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CONH}-$ hairpin linkage derived from γ -aminobutyric acid or a chiral hairpin linkage derived from 2,4-diaminobutyric acid;

X_1/X'_1 , X_2/X'_2 , X_3/X'_3 , X_4/X'_4 , X_5/X'_5 , X'_1/X'_1 , X'_2/X'_2 , X'_3/X'_3 , X'_4/X'_4 , and X'_5/X'_5 represent carboxamide binding pairs which bind DNA base pairs and are selected from the group consisting of H/P , P/H , P/H_m , I_m/P , and P/P , to correspond to the DNA base pair in the minor groove to be bound;

L represents an amino acid linking group selected from the group consisting of β -alanine and 5-aminovaleric acid (δ);

P represents a polyamide selected from the group consisting of $X_1X_2X_3X_4X_5X_6$,

$X_1X_2X_3X_4X_5X_6X_7X_8$, $X_1X_2X_3X_4X_5X_6X_7X_8X_9X_{10}$, and $X_1X_2X_3X_4X_5X_6X_7X_8X_9X_{10}X_{11}X_{12}$,

where X_1 - X_{12} are independently selected from the group consisting of β -alanine, pyrrole, hydroxypyrrole and imidazole; and

A represents a positive patch consisting of a rigid group adjacent to a positively charged group.

A represents a positive patch consisting of a rigid group adjacent to a positively charged group [such that a positive charge is delivered to the phosphate backbone or the major groove of a DNA molecule].

Art Unit: 1635

24. (Amended) The polyamide of claim 1 selected the group consisting of:

ImPyPyPy-γ-PyPyPyPy-β-RPR;
 ImImPyPy-γ-ImPyPyPy-β-RPR;
 ImPyPyPy-γ-PyPyPyPy-β-RPRRR;
 ImImPyPy-γ-ImPyPyPy-β-RPRRR;
 ImPyPyPy-γ-PyPyPyPy-β-R;
 ImPyPyPy-γ-PyPyPyPy-β-RP;
 ImPyPyPy-γ-PyPyPyPy-β-RGR;
 ImPyPyPy-γ-PyPyPyPy-β-R^DPR;
 ImPyPyPy-γ-PyPyPyPy-β-APR;
 ImPyPyPy-γ-PyPyPyPy-β-KPR;
 ImPyPyPy-γ-PyPyPyPy-β-RPK;
 ImPyPyPy-γ-PyPyPyPy-β-C7-RPR; and

the pharmaceutically acceptable salts thereof.

25. (Amended) A method of inhibiting gene expression comprising contacting a regulatory sequence of a gene with the polyamide of claim 1.

26. (Amended) A method of inhibiting gene expression comprising contacting a DNA molecule with the polyamide of claim 1 whereby the DNA molecule is conformationally constrained.